

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

EXPERIENTIA



Revue mensuelle des sciences pures et appliquées · Month
Monatsschrift für die Gebiete der Naturwissenschaft · Еже-

Rivista mensile di scienze pure e applicate · Revista r

ores: H. Aebi, Bern · D. Arigoni, Zürich · P. S. Chen, Zürich · A. Ple

Redactor: H. Mislin, Basel ·

Birkhäuser Verlag, Basel · Boston

GEORGE WASHINGTON UNIVERSITY
PAUL HINGELFORD HEALTH SCIENCES LIBRARY
2300 EYE STREET, N. W.
WASHINGTON, D. C. 20037

AUG 6 1980

Table des matières · Inhaltsverzeichnis · Sommario · Contents · Содержание · Índice

THE NEWMAN LIBRARY INC. 11 MARSHFIELD AVENUE

Oralia

carbon dioxide problem. An interdisciplinary survey

Introduction

H. Bach 767

The carbon cycle: sources and sinks of atmospheric CO₂

H. Kohlmaier, U. Fischbach, G. Kratz, H. Bröhl and H. Schunck 769

The effect of the atmosphere-biosphere exchange on the global carbon cycle

H. Hampicke 776

Past and future emission of CO₂

R. M. Rotty 781

Prediction of future CO₂ concentrations in the atmosphere

C. Siegenthaler and H. Oeschger 783

Physiological aspects of increased CO₂ concentration

H. van Keulen, H. H. van Laar, W. Louwerse and J. Goudriaan 786

The impact of different energy options on atmospheric CO₂ levels

E. Niehaus 792

Climatic effects of increasing atmospheric CO₂ levels

H. Bach 796

Measures of CO₂ control

C. Marchetti 806

Conclusions

H. Oeschger 808

Oralia

Mathematica, Physica

The critical rotation of gravitating degenerate cylinders

Z. F. Seidov 813

Biochimica

Endin, a C₂₂ degraded terpene from the sponge *Phyllospongia*

R. Kazlauskas, P. T. Murphy and R. J. Wells 814

Synthesis of koenigicine

R. B. Sharma, R. Seth-Verma and R. S. Kapil 815

2-dimensional electrophoresis: A combination of crossed immunoelectrophoresis and dodecyl sulfate polyacrylamide gel electrophoresis for protein analysis

I. Yamamoto and Y. Tanada 816

amine oxidase inhibitors and histamine metabolism

M. Strolin Benedetti, J. F. Ancher and N. Sontag 818

The effect of pH and methylation on the interaction of deoxycho-

lesterol with rat liver alcohol dehydrogenase

M. Antonetta and G. M. Hanozet 820

omerization of dopa and cysteinyl-dopa in melanogenesis in

S. Eo, E. Novellino, F. Chioccare, G. Misuraca and G. Prota 822

The response of cyclic 3',5'-AMP and cyclic 3',5'-GMP phospho-

diesterases to experimental diabetes

B. Perez de Gracia, A. R. Rhoads and W. L. West 824

A comparison of 3 reagents in converting thiamine to thiochrome

in the presence of plant extracts and polyphenols

K. Ratanaulchai, S. Pikulkarntaleri and B. Panijpan 825

Biologica

Karyotypes of several frogs from Korea, Taiwan and the Philip-

pines

M. Kuramoto 826

Effect of ethanol and isopropanol on the activity of alcohol

dehydrogenase, viability and life-span in *Drosophila melanogaster*

and *Drosophila funebris*

L. Vilageliu Arqués and R. Gonzalez Duarte 828

Spectral sensitivity in a fresh-water Gastrotrich (*Lepidodermella*

squamatum Dujardin)

M. Balsamo 830

Identification of intracellular particles of pesticides in ciliate proto-

zoa by Raman microprobe

D. Dive, J. M. Devynck, G. Leroy, M. N. Fourmaux and Y. Moschetto 832

Neurosecretory system and storage of paraldehyde fuchsin positive

neurosecretory material in the dorsal aorta in an earwig, *Anisolabis*

annulipes Lucas (Dermaptera: Labiduridae)

Y. N. Singh and R. Narain 834

The effect of RNA synthesis inhibitors on prenucleolar bodies

Gloria Morcillo and C. de la Torre 836

Satellite DNA sequences and reproductive isolation in the *Droso-*

phila willistoni group

G. Corneo, L. Ceccherini Nelli, Donatella Meazza and F. J. Ayala 837

Absorption of glycine and proline from the small intestine of rats

infected with *Eimeria nieschulzi*

S. J. Ball, Christine E. Heading and B. Tranter 839

Physiologica

Effects of altered thyroid state on the inhibition produced by locus

coeruleus in Purkinje cells in the rat

Dolores Fuenmayor and J. A. Gonzalez-Vegas 841

Microsphere measurement of myocardial capillary bed in hypoxic

rats

A. J. Stere, N. W. Brister and A. Anthony 842

Catalase activity in electrically stimulated muscle

R. R. Jenkins and D. Newsham 843

Effects of inhibition and stimulation of Na⁺-K⁺ active transport on

the resting membrane input conductance of the guinea-pig ven-

tricle

T. M. Nosek 844

Dietary utilization of aliphatic alcohols by *Drosophila*

J. Van Herrewege, J. R. David and R. Grantham 846

vide paginam aversam

- Time course of excitatory and inhibitory states of bulbar respiratory modulated neurons
M. Falleri, H. R. O. Dinse and G. Böhrer 848
- Effect of sympathectomy on platelet aggregation and blood coagulation in rats
A. Mukhammedov and A. V. Suvorov 849
- Intraventricular administration of isoproterenol inhibits both heat production and heat loss mechanisms in rats
M. T. Lin, A. Chandra, Y. C. Fan and R. Sun 851
- Glutamate decarboxylase activity in brain regions of differentially-housed mice: a difference in the olfactory bulb
J. M. Blindermann, F. V. DeFeudis, M. Maitre, R. Misslin and P. Mandel 853

Pharmacologica

- Hypothermia induced in mice by enzyme-mediated depletion of serotonin
J. Roberts and H. J. Rosenfeld 854
- Multiple esterase forms in isolated hepatocytes and Kupffer cells of partially hepatectomized rats
D. N. Mayanski, V. I. Scherbakoff and Alevtina U. Grishanova 856
- The infectivity of polyhedra of nuclear polyhedrosis virus (N.P.V.) after passage through gut of an insect-predator
A. G. B. Beekman 858
- The adequacy of thiamine in liquid diets used in animal models of alcoholism
A. W. K. Chan, Donna L. Schanley and Florence W. Leong 859
- Effect of praseodymium on drug metabolism in rat liver smooth and rough endoplasmic reticulum
P. Arvela, B. v. Lehmann, O. Grajewski and E. Oberdisse 860
- Inhibition of uridinediphosphate glucuronyltransferase caused by furosemide
F. Sörgel, F. E. Beyhl and E. Mutschler 861
- Autoradiographic evidence for binding of ³H-flunitrazepam (Rohypnol®) to melanin granules in the cat eye
H. Kuhn 863
- Ureteral contractions induced by rat urine in vitro: probable involvement of renal kallikrein
M. Marin-Grez, G. Bönner and F. Gross 865
- Neuromuscular blocking activity of dibekacin, a new semisynthetic aminoglycoside antibiotic
A. G. Paradis, J. S. Tsouras, P. Latsoudes, T. Demetriou and C. Triantaphyllidis 867
- Myotoxic phospholipases A from snake venom, *Pseudechis colleti*, producing myoglobinuria in mice
D. Mehs and Y. Samejima 868

Anatomica, Histologica, Embryologica, Carcinologica

- The induction of chromosome aberrations in human lymphocytes by negativ π -mesons under conditions of anoxia and oxygenation
J. S. Prosser and S. J. Prisman 869

- Normal osteoclast number and function in rat pups lacking parathyroid hormone
M. Krukowski and A. J. Kahn 87
- Radioprotective action of haematoporphyrin on X-irradiated tryptophan
V. Pistoia and Carmela Murolo 87

Haematologica, Immunologica

- The effect of dietary administration of aspartic acid on thymus weight in C57 black mice
Iva Pipalová and M. Pospíšil 87
- An improved electrophoretic method for a screening program for haemoglobinopathies
P. C. Naoum, M. C. R. Moura Campos, M. F. Parenti and A. M. Szymanski 87
- Chromatin organization within nuclear blebs in leukocytes of *Xenopus laevis*
N. Chegini and N. Maclean 87

Endocrinologica

- Estradiol treatment reduces a cytosol androgen binding protein in male rat liver
N. Sato, M. Ota and S. Takahashi 87
- Evidence that a 'critical period' is involved in early estrogen-induced LH- and FSH-release during the estrous cycle in the rat
J. M. Geiger and Cl. Aron 87
- Binding of iproniazid to the polymeric forms of iodide peroxidase
A. Nagasaka, Leslie J. DeGroot and H. Hidaka 88
- Effect of repeated electroconvulsive shock on plasma noradrenaline and adrenaline in man
I. Klimes, R. Kvetňanský and M. Chmelová 88
- Molting processes and hormonal control: an in vitro model
J. P. Caruelle 88

Pro experimentis

- FITC-dextran as fluorescence and electron microscopic tracers in studies on capillary and cell permeability of the CNS
U. Heinzmann 88
- A simple method for the verification of a successful cannulation of the rat cerebral ventricles
I. Paakkari 88
- Immunogenicity of agarose-immobilized immune complexes
P. J. Higgins 88

Congressus

Instructions to authors

the damage of the periventricular structures. An additional advantage of this method is the simplicity and unexpensiveness of the technical tools as compared to the methods using pressure transducers and recording devices^{8,9}.

- 1 This work was done under a contract with the Association of Finnish Life Insurance Companies.
- 2 E. U. Gerdes, G. Klein, M. Nadjmi and G. Schaltenbrand, *J. Neurol.* 210, 183 (1975).

- 3 K. I. Melville and H. E. Shister, *Am. Heart J.* 53, 425 (1957).
- 4 L. Eriksson, *Acta physiol. scand.*, suppl. 444, 8 (1976).
- 5 H. Schmitt and H. Schmitt, *Eur. J. Pharmac.* 6, 8 (1969).
- 6 V. S. Vassiliev and A. O. Ramos, *Pharmacology* 13, 97 (1975).
- 7 E. K. Walls and T. B. Wishart, *Physiol. Behav.* 19, 171 (1977).
- 8 A. Hernandez, S. Ruiz and C. Paele, *Experientia* 31, 1475 (1975).
- 9 G. A. Goodrich, B. Greehey, T. B. Miller and J. R. Pappenheimer, *J. appl. Physiol.* 26, 137 (1969).

Immunogenicity of agarose-immobilized immune complexes¹

P. J. Higgins

Laboratory of Cell Biochemistry, Memorial Sloan-Kettering Cancer Center, 1275 York Avenue, New York (N.Y. 10021, USA), 8 October 1979

Summary. A novel method is described for production of heterologous antisera to a specific tumor-associated murine antigen by immunization with agarose-trapped immune complexes.

Preparation of high titer heterologous antisera to specific tumor-associated antigens is complicated by the use of crude tissue extracts as immunogen in which the relative concentration of the antigen is quite low. Where the tumor-associated antigen is present in moderate quantities, reasonable quality antisera can be obtained by immunization with unfractionated tissue extracts and appropriate serum absorptions. Such antisera have been used to precipitate antigen from solution and the immune complexes, in turn, employed as immunogens to elicit the formation of a more highly specific antiserum². The limiting factors, however, of sufficiently absorbed precipitating antibody, low concentrations of specific antigen in crude tissue extracts and difficulties in the efficient processing of minute precipitates combine to make effective immunization with solution-precipitated immune complexes a prodigious task.

A simple method for the preparation of high titer antisera to a specific tumor-associated antigen, murine γ -FA³⁻⁵, is described. This procedure should be applicable to studies of diverse tissue antigens.

Methods and results. Immunization of rabbits with saline extracts of a mouse fibrosarcoma, antiserum absorption and subsequent identification of an antigen common to tumor, fetal and adult splenic tissue, termed γ -FA, has been described³⁻⁵. Radial immunodiffusion plates⁶ contained 0.1 ml of absorbed anti- γ -FA serum and 2.5 ml of 1% agarose (w/v) in Beckman B-2 buffer, pH 8.6. Antigen wells (3.7 mm in diameter) were cut into the agarose gel, filled with \sim μ l of the 10,000 \times g supernatant fraction of a

10^{-3} M Tris, pH 7.5, homogenate of normal adult mouse spleen and the plates incubated at 37°C for 72 h to allow for precipitin ring formation. The agarose slabs were then dialyzed with stirring against daily 200-ml changes of phosphate-buffered saline (PBS) for 3 weeks at 4°C in order to remove unbound protein. The use of a 1% agarose gel facilitated this removal while antigen-antibody complexes remained trapped within the gel matrix. After dialysis, sections of gel containing precipitin rings were cut out, passed several times through an 18-gauge needle and frozen at -20°C in twice the volume of PBS. On day 1, 14 and 21, 1 ml of agarose-immobilized immune complexes was emulsified in 1 ml of complete Freund's adjuvant and inoculated s.c. into a New Zealand white rabbit (on day 21 incomplete adjuvant was substituted for complete adjuvant). 22 days later the rabbit was bled and 1-ml aliquots of the antiserum inoculated i.p. into each of several adult C57 mice for in vivo absorption. After 24 h, the mice were bled and the antibody activity of the absorbed antiserum compared with that of the original anti- γ -FA serum. Clearly, the antiserum to γ -FA-anti- γ -FA immune complexes possessed all the precipitin specificity of the original anti- γ -FA serum but at a much higher titer (table) and, unlike the original antiserum, yielded a positive indirect immunofluorescence test⁴ on methanol-fixed rat hepatoma cells (figure). This antigen was previously thought to be synthesized only by in vivo propagated tumor cells⁴ and the present data, therefore, provide the 1st direct evidence for production of γ -FA by transformed cells. Moreover, reten-




Fig. 1. Indirect immunofluorescence test of in vitro propagated rat tumor cells isolated from a transplanted hepatoma⁷. A Original anti- γ -FA serum (1:20); B anti- γ -FA immune complexes (1:60). UV light microscopy. BG12-53/44 filters, Zeiss Photomicroscope.

Table 1. Comparative precipitin tests^a

Test antigen	Original anti- γ -FA (undiluted)	Anti- γ -FA immune complexes (1:30) ^b
Meth A serum ^c	+	+++
Meth A tumor ^d	+	++++
Normal serum ^e	-	-
Adult spleen ^f	+	++++
Viscera ^g	-	-

^a Hyland Immuno-Plates, pattern 'D'. Preliminary experiments indicated the precipitin line which formed upon interaction of the original anti- γ -FA serum with a saline extract of adult mouse spleen and that which formed by interaction of anti- γ -FA immune complexes with the same splenic tissue extract was one of identity.

^b Calculated dilution after in vivo absorption. ^{c,d} Serum and saline extract of tumor tissue obtained from a mouse bearing a transplanted 3-methylcholanthrene-induced fibrosarcoma. ^{e,f} Obtained from normal adult mice. ^g Individual saline extracts of adult mouse liver, kidney, brain, heart, lung, testes and small intestine pooled from several mice.

tion of antibody activity after in vivo absorption in mice (which have γ -FA-positive spleens) suggests that γ -FA is not a cell surface antigen. Preparation of a high titer anti- γ -FA serum by immunization with agarose-trapped immune complexes will facilitate future in vitro studies of various aspects of cellular transformation. In addition, the method described should be adaptable to diverse tissue antigen systems.

- Supported in part by a Biomedical Research Support Grant and Grant RO1-CA-20019-04 from the National Cancer Institute.
- S. Nishi, *Cancer Res.* 30, 2507 (1970).
- C. Tong, E. H. Stonehill, P. J. Higgins and A. Bendich, *Eur. J. Cancer* 14, 147 (1978).
- P. J. Higgins, C. Tong, E. Borenfreund and A. Bendich, *Eur. J. Cancer* 15, 423 (1979).
- P. J. Higgins, C. Tong, E. Borenfreund, R. S. Okin and A. Bendich, *Br. J. Cancer* 40, 253 (1979).
- G. Mancini, A. O. Carbonara and J. F. Heremans, *Immunochemistry* 2, 235 (1965).
- E. Borenfreund and P. J. Higgins, *Ann. N.Y. Acad. Sci.*, in press (1980).

CONGRESSUS

Great Britain

4th meeting of European neuroscience association

Brighton, 13-20 September 1980

Plenary lectures will be presented by H. Barlow, M. Burger, A. Burgen, E. Zaidel, S. Thesleff, P. Andersen, D. Ploog and L. Heimer.

For further information, please write to the secretary of the meeting, 142 Oxford Road, Cowley, Oxford OX4 2DZ, England.

Switzerland

12th FEMS symposium "Microbial degradation of xenobiotics and recalcitrant compounds"

Zürich, 15-17 September 1980

Main lectures on "Xenobiotics in the environment", "Cellular and subcellular aspects of biodegradation research" and "Biodegradation of specific compounds". For information contact Prof. Th. Leisinger, Mikrobiologisches Institut ETH, ETH-Zentrum, CH-8092 Zürich.

Instructions to authors

Experientia is published on the 15th of every month and can be obtained in any country through booksellers or from the publishers. All communications to the editors should be addressed to the publishers. All manuscripts for publication in a given number must be in the hands of the editors 3 months before publication.

Articles of general scientific interest, of interdisciplinary character: briefly stated and hitherto unpublished original reports of sufficient novelty value.

Experiments Papers in which animal experiments have been conducted without using the appropriate anaesthesia will not be accepted.

Manuscripts (including all figures and tables) should be submitted in duplicate.

Text should not exceed 2-3 typewritten pages (50-60 lines), 1-2 relevant figures or tables. English summary of maximum 4 lines. Abbreviations should be properly explained. References should be numbered consecutively and be presented on a separate page. Name and address have to be placed directly under the title. Linguistically inadequate manuscripts will be returned. Manuscripts in languages other than English should be supplemented by an English translation of the title. Footnotes should be avoided.

Figures Illustrations should be separate from the text, with the author's name on the back in soft pencil. The desired labelling should be shown on a second set of figures, which will be used as a model for

inscriptions. Drawings for reproductions should be on good paper in Indian ink, photographs should be supplied as glossy positive prints. The illustrations should be at least one and a half times as large as the definitive size desired. Over-large figures can be easily damaged in the mail. Captions should be self-explanatory, without reference to the text.

Tables should be provided with a title and with self-explanatory captions.

Headings In submitting their manuscript to *Experientia*, authors are requested to indicate one of the headings mentioned below, under which they would wish to place their short communication.

1. Mathematics and Physics; 2. Cosmology, Astronautics, Cosmonautics; 3. Mineralogy, Geophysics, Oceanography; 4. Inorganic and Physical Chemistry; 5. Organic Chemistry; 6. Biophysics; 7. Molecular Biology, Cellular Biology; 8. Genetics; 9. Botany; 10. Zoology; 11. Ecology; 12. Biochemistry (analytic and synthetic); 13. Biochemistry (Enzymes, Metabolism); 14. Physiology; 15. Neurology; 16. Pharmacology, Toxicology, Pathology; 17. Experimental Gerontology; 18. Anatomy, Histology, Cytology, Histochemistry; 19. Embryology; 20. Endocrinology; 21. Circulation, Cardiology, Angiology; 22. Nutrition, Gastroenterology; 23. Hematology, Serology; 24. Immunology, Allergy; 25. Microbiology, Parasitology, Chemical Therapeutics; 26. Oncology, Carcinology, Cytostatics; 27. Radiology.

Reprints The authors receive 50 reprints, without cover, free of charge. Price-list for further reprints is available.